



# Sustained rapid shrinkage of Yukon glaciers since the 1957–1958 International Geophysical Year

N. E. Barrand<sup>1</sup> and M. J. Sharp<sup>1</sup>

Received 3 December 2009; revised 14 January 2010; accepted 12 February 2010; published 1 April 2010.

[1] Glaciers in the Yukon, NW Canada, lost 22% of their surface area during the 50 years following the 1957–58 International Geophysical Year, coincident with increases in average winter and summer air temperatures and decreases in winter precipitation. Scaling these results to ice volume change, we obtain a total mass loss of  $406 \pm 177$  Gt, which accounts for  $1.13 \pm 0.49$  mm of global sea-level rise. Yukon glaciers thinned by  $0.78 \pm 0.34$  m yr<sup>-1</sup> water equivalent, a regional thinning rate exceeded only by mountain glaciers in Patagonia and Alaska. Our scaling analysis suggests the remaining glaciers have the capacity to contribute a further 5.04 mm to global sea-level rise. **Citation:** Barrand, N. E., and M. J. Sharp (2010), Sustained rapid shrinkage of Yukon glaciers since the 1957–1958 International Geophysical Year, *Geophys. Res. Lett.*, *37*, L07501, doi:10.1029/2009GL042030.

## 1. Introduction

[2] The response of glaciers and ice caps (GIC) to climate change is an issue of significant public and scientific concern because of its implications for global sea-level rise (SLR). In addition to ocean thermal expansion, the present day sea-level budget attributes ~half of observed SLR to the ice sheets and ~half to mass losses from GIC [Cazenave *et al.*, 2009]. The GIC contribution to SLR has accelerated over the past decade [Meier *et al.*, 2007], yet the errors associated with estimates of this contribution are often large. Reducing them requires comprehensive studies of regional-scale glacier changes, particularly from undersampled areas that have experienced substantial climate change in recent decades.

[3] Although ~10,000 km<sup>2</sup> of the Yukon, NW Canada, is glacier covered (Figure 1), very little is known about the health of these glaciers due to an absence of field measurements and the difficulty of estimating regional-scale volume changes. NW North America has experienced significant climate change over the past half century, with increases in average winter (October–April,  $2.0 \pm 0.8^\circ\text{C}$ ) and summer (May–September,  $1.0 \pm 0.4^\circ\text{C}$ ) air temperatures and a small increase in precipitation (albeit significant at just 17% of sites) recorded at 67 National Oceanic and Atmospheric Administration (NOAA) and Environment Canada weather stations [Arendt *et al.*, 2009]. Of these stations, the 14 located within the Yukon record similar summer warming ( $0.99^\circ\text{C}$ ), yet a reduction in winter precipitation (on average,  $-22$  mm). Regional climate warming, reduction in

winter precipitation and increases in freezing level heights (FLHs) [Arendt *et al.*, 2009] suggest that both the maritime (St. Elias (Figure 1b)) and interior (Mackenzie (Figure 1c)) glacier populations of the Yukon may have experienced large losses during the past 50 years.

[4] Existing measurements of glacier mass loss from the St. Elias mountains are limited either spatially (extrapolation of centreline thinning rates from just three sites, of which only one, Kaskawulsh glacier, is located entirely within the Yukon [Arendt *et al.*, 2002]) or temporally (GRACE satellite gravity solutions from 2003–2007 [Luthcke *et al.*, 2008]). The latter study estimated thinning rates of  $0.63 \pm 0.09$  m yr<sup>-1</sup> water equivalent (w.e.) during this recent time period. No study to date has measured regional glacier changes in the Mackenzie mountains. Here we present the first comprehensive multi-decadal assessment of Yukon glacier changes (including both St. Elias and Mackenzie glacier populations) and their associated contribution to SLR, and evaluate the potential SLR contribution from the remaining glaciers.

## 2. Methods


### 2.1. Study Area and Data Sources

[5] We investigated area changes of all 1396 glaciers in the Yukon during the half century between the first International Geophysical Year (IGY1, 1957–58) and the third International Polar Year (IPY3, 2007–09). Our study area comprises the regions between 60–62°N, 135–141°W (St. Elias mountains), and 63–65°N, 130–133°W (Mackenzie mountains) (Figure 1). We delineated glacier boundaries from 802 individual 1:50,000-scale aerial photographs covering every glacier in the Yukon from late summer 1958–60 acquisitions by the Canadian Land Survey, and from 11 Landsat-7 Enhanced Thematic Mapper (ETM+) satellite images acquired in summers 2006–08. Aerial photography prints were digitally scanned at 500 dpi (~2 m ground resolution) and spatially referenced to the orthorectified satellite imagery using >5 ground control points per photograph.

### 2.2. Derivation of Area Changes

[6] Glacier margins were manually digitized as polygon shapefiles (surface area = margin minus bedrock, nunataks). Glaciers flowing from drainage divides were digitized separately based on ice-surface contours from Canadian National Topographic Series (NTS) digital maps. Margins of debris-covered glaciers were delineated using map contours and visual interpretation. Polygons were delineated in historical aerial photographs, overlaid onto contemporary satellite imagery and edited where change had occurred. Margins were edited only where change was clearly evident,

<sup>1</sup>Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, Alberta, Canada.



**Figure 1.** (a) Location map showing the extent of glacial ice (black shading) in the Yukon, NW Canada. Insets display detail of (b) St Elias and (c) Mackenzie Mountain glacier populations. Background image is a shaded relief mosaic derived from GTOPO/SRTM digital elevation data.

an approach which may conservatively estimate retreat at debris-covered glaciers. We included surge-type glaciers in our area calculations, having identified 19 surging glaciers (totaling 924.2 km<sup>2</sup>, or 10% of the total area) from morphological evidence and previously published material [e.g., Clarke and Collins, 1984; Clarke *et al.*, 1986]. Due to limited photographic coverage glaciers overlying international and provincial borders were cropped along administrative boundary lines. Area change errors were calculated by selecting a random sample of 50 glaciers and deriving an ice-area uncertainty fraction based on the root sum of squares (RSS) of individual error components appropriate to each glacier ( $\pm 15$  m for clear-ice margins,  $\pm 45$  m for debris/snow-covered margins,  $\pm 15$  m for image registration uncertainty). The fractions appropriate to each of these 50 glaciers were then averaged and scaled to the entire dataset [e.g., DeBeer and Sharp, 2007].

### 2.3. Volume/Area Scaling

[7] We calculated volumes and volume changes for each individual glacier (excluding surge-type glaciers) by scaling glacier area ( $A$ ) to ice volume ( $V$ ) using the power law relation:

$$V = c_a A^\gamma$$

[Chen and Ohmura, 1990; Bahr *et al.*, 1997], where  $c_a$  is equal to 0.28 m<sup>(3-2 $\gamma$ )</sup>, and  $\gamma$  is a scaling exponent equal to 1.375. These coefficients were calibrated from direct measurements of area and volume change of non-tidewater glaciers in the nearby Western Chugach Mountains, Alaska [Arendt *et al.*, 2006], and are therefore the most appropriate available for our (non-tidewater valley glacier) sample. While this approach may generate large errors at the individual glacier scale, it is suitable (errors typically <25%) for large, regional-scale glacier populations [Meier *et al.*, 2007]. Yukon IGY1 and IPY3 glacier inventory data and full area

and volume calculations are available for download as auxiliary material.<sup>1</sup>

[8] To give an idea of the sensitivity of volume estimates to the choice of scaling coefficients we calculated a range of estimates using coefficients derived from glaciers in the Alps, Cascades and similar regions [Chen and Ohmura, 1990], the Canadian Cordillera (south of our study area [DeBeer and Sharp, 2007]), and from solely physical considerations [Bahr *et al.*, 1997] (see Table 1). While individual volume estimates based on Chugach coefficients are preferred, use of other coefficients gives a range of values. As uncertainty in volume changes from the choice of scaling coefficients is typically an order of magnitude greater than those due to area measurement errors, we consider volume uncertainty due to area change errors as negligible. Instead we calculate the uncertainty of scaled volume changes as the maximum difference between estimates using Chugach coefficients and those using the different coefficients provided in Table 1.

[9] We acknowledge an additional uncertainty resulting from areas cropped along administrative boundary lines. Of the 1396 glaciers in our inventory, 52 were cropped at the Yukon-Alaska boundary. In 20–25 cases the cropping took place along a drainage divide (digitized separately, see above). While it is difficult to accurately assess the uncertainty of scaled volumes for the remaining cropped glaciers, it is likely that these volumes will be underestimated. As the number of glaciers affected by this issue is small (2% of the total sample size) we consider this uncertainty to be insignificant compared to scaling coefficient errors.

[10] We converted volume changes to mass changes by assuming a constant density-depth profile [Bader, 1954] (that is, considering volume losses to consist of glacier ice

<sup>1</sup>Auxiliary materials are available at <ftp://ftp.agu.org/apend/gl/2009gl042030>.

**Table 1.** Sensitivity of Yukon Glacier Mass Loss Estimates to Published Scaling Coefficients  $c_a$  and  $\gamma$  in the Power Law Relation,  $V = c_a A^\gamma$ 

$c_a/\gamma$	Coefficient Source	Mass Loss (Gt, 1958–60–2006–08)	Difference From <sup>a</sup> (Gt)
0.280 m <sup>(3-2<math>\gamma</math>)</sup> /1.375	<i>Arendt et al.</i> [2006]	406 <sup>a</sup>	
0.206 m <sup>(3-2<math>\gamma</math>)</sup> /1.360	<i>Chen and Ohmura</i> [1990]	229	177 (–44%)
0.013 m <sup>(3-2<math>\gamma</math>)</sup> /1.520	<i>Chen and Ohmura</i> [1990]	256	150 (–37%)
0.115 m <sup>(3-2<math>\gamma</math>)</sup> /1.405	<i>Chen and Ohmura</i> [1990]	285	121 (–30%)
0.210 m <sup>(3-2<math>\gamma</math>)</sup> /1.360	<i>DeBeer and Sharp</i> [2007]	233	173 (–43%)
0.191 m <sup>(3-2<math>\gamma</math>)</sup> /1.375	<i>Bahr et al.</i> [1997]	277	129 (–32%)

<sup>a</sup>Yukon mass loss estimate calculated using scaling coefficients calibrated from glaciers in the nearby Western Chugach Mountains [*Arendt et al.*, 2006]. The fourth column provides total and percentage differences from this value, as calculated using five different sets of scaling coefficients.

only) and multiplying by the ratio of the density of ice to water ( $d_i/d_w = 0.917$  [e.g., *Paterson*, 1994]). Thickness change rates were calculated by dividing mass changes by the average of the old and new glacier areas [e.g., *Echelmeyer et al.*, 1996; *Arendt et al.*, 2002].

### 3. Results

[11] During the 50 years between IGY1 and IPY3, total ice area in the Yukon shrank from 11,622 km<sup>2</sup> in 1958–60 to 9,081 km<sup>2</sup> in 2006–08, a loss of 2,541 ± 189 km<sup>2</sup>. This represents a 22% loss of glacier area over the course of 50 years. None of the surge-type glaciers we identified advanced during this time period. Viewing glacier area change (as a percentage of initial area) against initial area (1958–60) showed that glaciers of all sizes experienced substantial losses (Figure 2). In ten cases, glaciers >50 km<sup>2</sup> lost ≥20% of their 1958–60 area.

[12] Applying the volume/area scaling approach we calculated a total (w.e.) mass of Yukon glaciers during IGY1 (1958–60) as 2218 Gt. The equivalent mass in IPY3 (2006–08) was 1812 Gt. Our measured volume changes equate to mass losses in the range 155 ± 68 Gt (assuming all mass lost is old snow ( $d_i/d_w = 0.35$ )) to 406 ± 177 Gt (assuming all mass lost is glacier ice). Given that most of the losses result from glacier melt and occurred at low elevations on the glacier, a value towards the upper limit of this range seems likely. Assuming all mass lost is glacier ice, we calculate an annual loss rate of 8.12 ± 3.54 Gt yr<sup>-1</sup>. Averaging this mass loss over the area of the Earth's oceans ( $3.61 \times 10^{14}$  m<sup>2</sup> [e.g., *Bindoff et al.*, 2007]), we find that loss of Yukon glaciers contributed a total of 1.13 ± 0.49 mm to global SLR between IGY1 and IPY3. This corresponds to an annual SLR contribution of 0.02 ± 0.009 mm yr<sup>-1</sup> between 1958–60 and 2006–08. Our scaling approach suggests that the remaining mass of glacier ice in the Yukon is equivalent to a global SLR contribution of 5.04 mm.


### 4. Discussion

[13] The use of a volume/area scaling-based approach was necessary due to a lack of accurate topographic data from which to make direct measurements of volume change. Our findings may be considered regionally representative as we measured area changes of every glacier in the Yukon, inclusive of both the St Elias (Figure 1b) and Mackenzie Mountain ranges (Figure 1c). While we are confident that our choice of scaling coefficients was appropriate, a sensitivity analysis showed that the maximum difference between mass changes derived from Chugach mountain scaling


coefficients and those derived from previously published coefficients (based on physical considerations and empirical findings at glacier populations elsewhere) was ~±40% (Table 1). In the absence of independent validation data, we consider this value to represent a conservative estimate of measurement uncertainty.

[14] While an error bound of ~±40% was appropriate for our regional-scale volume changes, we were able to compare scaling results with direct measurements at a single site in the Yukon; Kaskawulsh glacier (~60°40'N, 138°50'W). Airborne laser altimetry data show that the glacier lost an average of 0.40 Gt yr<sup>-1</sup> from 1977–2005 (A. Arendt, personal communication, 2009). Our volume/area scaling results suggest that Kaskawulsh lost 0.61 Gt yr<sup>-1</sup> between IGY1 (1958–60) and IPY3 (2006–08). Although our mass loss is more negative than that derived from direct measurements, the similarity between these two values is encouraging, particularly given the different time interval, the lack of errors in the altimetry estimate, and the uncertainties of assuming all volume losses to consist of glacier ice and from interpreting volume scaling results from individual glaciers.

[15] We have shown that substantial changes in Yukon glacier area between IGY1 and IPY3 were coincident with a period of general summer warming and reduction in winter precipitation recorded at 14 Environment Canada weather stations. We also examined interannual temperature variability in the Yukon, from National Centre for Environ-



**Figure 2.** Plot of percentage glacier area change 1958–60–2006–08 as a function of initial (1958–60) glacier area. Nine glaciers larger than 200 km<sup>2</sup> are not shown.



**Figure 3.** Half yearly averaged 2 m air temperatures in the Yukon, NW Canada, (a) winter, October–March; (b) summer, April–September, from NCEP/NCAR reanalysis model output.

mental Prediction (NCEP)/National Centre for Atmospheric Research (NCAR) reanalysis model output [Kalnay *et al.*, 1996]. NCEP/NCAR temperatures are based strongly on observations and thus fall within the most reliable class of all NCEP variables. Mean air temperatures in the Yukon between 1950 and the present (2010) show a general winter warming trend since 1950 (Figure 3a), and summer warming since ~1970 (Figure 3b). Reduction in precipitation and summer warming result in increasingly negative glacier mass balances as a result of decreases in winter accumulation and increases in summer melting. Winter warming however, may also be important for glaciers in this region. Previous research has shown that (lower elevation maritime) glaciers in the St. Elias mountains have winter FLHs above 6–27% of their ablation areas [Arendt *et al.*, 2009], meaning that they melt during both summer and winter seasons. Rising winter temperatures will increase FLHs, exposing more glacier area to above freezing temperatures and increasing the proportion of winter precipitation that falls as rain. The magnitude of this effect will depend on both the amount of shift in FLH and the glacier's area-altitude distribution [Arendt *et al.*, 2009].

[16] The area-averaged thinning rate of Yukon glaciers between IGY1 and IPY3 was  $0.78 \pm 0.34 \text{ m yr}^{-1}$  w.e. This value is in agreement with an entirely independent estimate of St. Elias mass balance between April 2003 and March 2007 from GRACE satellite gravimetry (mascon region 7,  $0.63 \pm 0.09 \text{ m yr}^{-1}$  w.e. [Luthcke *et al.*, 2008]). The similarity of these two results indicates that recent rapid thinning

may have been sustained for much longer than previously assumed. As this value is area-averaged it is possible to compare it with published estimates of regional thinning rates for GIC populations elsewhere in the world. Our thinning rate for Yukon glaciers is larger (although within errors) than the  $0.52 \text{ m yr}^{-1}$  loss of Gulf of Alaskan glaciers between the mid 1950s and mid 1990s, yet smaller than the  $1.80 \text{ m yr}^{-1}$  thinning between the mid 1990s and early 2000s [Arendt *et al.*, 2002]. The discrepancy between these two values is likely a result of dynamic thinning of large tidewater glaciers in the Gulf of Alaska (none of which are present within the neighbouring Yukon). Our thinning rate is less than those of the Patagonian icefields [Rignot *et al.*, 2003; Chen *et al.*, 2007] (and mid 1990s–mid 2000s Alaska), yet greater than regional loss rates reported throughout the Canadian [Abdalati *et al.*, 2004] and Norwegian [Bamber *et al.*, 2005; Nuth *et al.*, 2010] Arctic archipelagos.

## 5. Conclusions

[17] Our calculation of mass loss (comprising measurements from every glacier) provides the first comprehensive multi-decadal estimate of the SLR contribution from Yukon glaciers and will help to reduce the uncertainty of GIC contributions to global SLR. Outside the ice sheets, the regional thinning rate we report here is exceeded only by Patagonian glaciers and by glaciers in Alaska during the period from the mid 1990s to the mid 2000s. Our findings indicate that Yukon glaciers are sensitive to climate perturbations and have experienced significant losses in response to local climate forcing. Our scaling analysis suggests that the remaining glacial ice in the Yukon would contribute a further 5.04 mm to global SLR were it to melt completely.

[18] **Acknowledgments.** Canadian Land Survey aerial photographs were made available from the archive of the National Hydrological Research Institute, Saskatoon, SK, and Landsat 7 ETM+ scenes were obtained from the United States Geological Survey (<http://landsat.usgs.gov>). This work was funded by grants from the Canadian Federal International Polar Year program and the Natural Sciences and Engineering Research Council. We thank Patrick O'Callaghan for digitizing assistance and Anthony Arendt, Nancy Baron, Alberto Reyes, and two anonymous reviewers for comments which improved the manuscript.

## References

- Abdalati, W., W. Krabill, E. Frederick, S. Manizade, C. Martin, J. Sonntag, R. Swift, R. Thomas, J. Yungel, and R. Koerner (2004), Elevation changes of ice caps in the Canadian Arctic Archipelago, *J. Geophys. Res.*, *109*, F04007, doi:10.1029/2003JF000045.
- Arendt, A., K. Echelmeyer, W. Harrison, C. Lingle, and V. Valentine (2002), Rapid wastage of Alaska glaciers and their contribution to rising sea level, *Science*, *297*, 382–386.
- Arendt, A., K. Echelmeyer, W. Harrison, C. Lingle, S. Zirnheld, V. Valentine, B. Ritchie, and M. Druckenmiller (2006), Updated estimates of glacier volume changes in the western Chugach Mountains, Alaska, and a comparison of regional extrapolation methods, *J. Geophys. Res.*, *111*, F03019, doi:10.1029/2005JF000436.
- Arendt, A., J. Walsh, and W. Harrison (2009), Changes of glaciers and climate in northwestern North America during the late twentieth century, *J. Clim.*, *22*, 4117–4134.
- Bader, H. (1954), Sorge's law of densification of snow on high polar glaciers, *J. Glaciol.*, *2*(15), 319–322.
- Bahr, D., M. Dyurgerov, and M. Meier (1997), The physical basis of glacier volume-area scaling, *J. Geophys. Res.*, *102*(20), 355–362.
- Bamber, J., W. Krabill, V. Raper, J. Dowdeswell, and J. Oerlemans (2005), Elevation changes measured on Svalbard glaciers and ice caps from airborne laser data, *Ann. Glaciol.*, *42*, 202–208.
- Bindoff, N., *et al.* (2007), Observations: Oceanic climate change and sea level, in *Climate Change 2007: The Physical Science Basis. Contribu-*

- tion of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by S. Solomon et al., pp. 385–432, Cambridge Univ. Press, Cambridge, U. K.
- Cazenave, A., K. Dominh, S. Guinehut, E. Berthier, W. Llovel, G. Ramillien, M. Ablain, and G. Larnicol (2009), Sea level budget over 2003–2008: A reevaluation from GRACE space gravimetry, satellite altimetry and Argo, *Global Planet. Change*, *65*, 83–88.
- Chen, J., and A. Ohmura (1990), Estimation of alpine glacier water resources and their change since the 1870s, in *Hydrology in Mountainous Regions I: Hydrological Measurements, The Water Cycle, Lausanne Symposia*, edited by H. Lang and A. Musy, *LAHS Publ.*, *193*, 127–135.
- Chen, J. L., C. R. Wilson, B. D. Tapley, D. D. Blankenship, and E. R. Ivins (2007), Patagonia Icefield melting observed by Gravity Recovery and Climate Experiment (GRACE), *Geophys. Res. Lett.*, *34*, L22501, doi:10.1029/2007GL031871.
- Clarke, G., and S. Collins (1984), The 1981–1982 surge of Hazard glacier, Yukon-Territory, *Can. J. Earth Sci.*, *21*(3), 297–304.
- Clarke, G., J. Schmok, C. Ommanney, and S. Collins (1986), Characteristics of surge-type glaciers, *J. Geophys. Res.*, *91*(B7), 7165–7180.
- DeBeer, C., and M. Sharp (2007), Recent changes in glacier area and volume within the southern Canadian Cordillera, *Ann. Glaciol.*, *46*, 215–221.
- Echelmeyer, K., W. Harrison, C. Larson, J. Mitchell, J. DeMallie, B. Rabus, G. Adalgeirsdóttir, and L. Sombadier (1996), Airborne surface profiling of glaciers: A case study in Alaska, *J. Glaciol.*, *42*(142), 538–547.
- Kalnay, E., et al. (1996), The NCEP/NCAR 40-year reanalysis project, *Bull. Am. Meteorol. Soc.*, *77*(3), 437–471.
- Luthcke, S., A. Arendt, D. Rowlands, J. McCarthy, and C. Larsen (2008), Recent glacier mass changes in the Gulf of Alaska region from GRACE mascon solutions, *J. Glaciol.*, *54*(188), 767–777.
- Meier, M., M. Dyurgerov, U. Rick, S. O’Neel, W. Pfeffer, R. Anderson, S. Anderson, and A. Glazovsky (2007), Glaciers dominate eustatic sea-level rise in the 21st century, *Science*, *317*(5841), 1064–1067.
- Nuth, C., G. Moholdt, J. Kohler, J. Hagen, and A. Kääb (2010), Svalbard glacier elevation changes and contribution to sea level rise, *J. Geophys. Res.*, *115*, F01008, doi:10.1029/2008JF001223.
- Paterson, W. (1994), *The Physics of Glaciers*, 3rd ed., Pergamon, Oxford, U. K.
- Rignot, E., A. Rivera, and G. Casassa (2003), Contribution of the Patagonian icefields of South America to sea level rise, *Science*, *302*, 434–437.

---

N. E. Barrand and M. J. Sharp, Department of Earth and Atmospheric Sciences, University of Alberta, 1-26 Earth Sciences Bldg., Edmonton, AB T5K 1J8, Canada. (barrand@ualberta.ca)